

(R)-1-Phenylethanaminium (S)-4-chloro-mandelate

Quan He,^a Michael C. Jennings,^{b*} Sohrab Rohani,^c Jesse Zhu^c and Hassan Gomaa^c

^aDepartment of Chemical Engineering, Ningbo University of Technology, Ningbo 315016, People's Republic of China, ^bDepartment of Chemistry, University of Western Ontario, London, Ontario, N6A 5B7, Canada, and ^cDepartment of Chemical and Biochemical Engineering, University of Western Ontario, London, Ontario, N6A 5B9, Canada

Correspondence e-mail: mjenning@uwo.ca


Received 17 January 2008; accepted 31 January 2008

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.003 \text{ \AA}$; R factor = 0.045; wR factor = 0.116; data-to-parameter ratio = 17.9.

The absolute configuration of the title complex, $C_8H_{12}N^+ \cdots C_8H_6ClO_3^-$ or $[R\text{-}C_6H_5C(H)CH_3NH_3][S\text{-}4\text{-Cl}C_6H_4C(H)\text{-}(OH)CO_2]$, has been confirmed by the structure determination. In the crystal structure, intermolecular O—H \cdots O and N—H \cdots O hydrogen bonds form a two-dimensional network perpendicular to the c axis.

Related literature

For background information and the crystal structure of the *R,R* diastereomer of the title compound, see: He *et al.* (2007).



Experimental

Crystal data

$C_8H_{12}N^+ \cdot C_8H_6ClO_3^-$
 $M_r = 307.76$
Monoclinic, $P2_1$
 $a = 10.4091 (7) \text{ \AA}$

$b = 5.7635 (4) \text{ \AA}$
 $c = 13.2544 (10) \text{ \AA}$
 $\beta = 96.831 (4)^\circ$
 $V = 789.52 (10) \text{ \AA}^3$

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.25 \text{ mm}^{-1}$

$T = 150 (2) \text{ K}$
 $0.45 \times 0.15 \times 0.08 \text{ mm}$

Data collection

Nonius KappaCCD diffractometer
Absorption correction: multi-scan (*SORTAV*; Blessing, 1995)
 $T_{\min} = 0.787$, $T_{\max} = 0.984$
7943 measured reflections
3431 independent reflections
2881 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.072$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.115$
 $S = 1.05$
3431 reflections
192 parameters
1 restraint

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
1436 Friedel pairs
Flack parameter: -0.03 (8)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O9—H9A \cdots O9 ⁱ	0.84	2.26	2.939 (2)	139
O9—H9A \cdots O11 ⁱ	0.84	2.09	2.826 (2)	146
N13—H13A \cdots O12 ⁱⁱ	0.91	1.89	2.798 (2)	172
N13—H13B \cdots O11 ⁱⁱ	0.91	1.83	2.731 (2)	169
N13—H13C \cdots O12 ⁱⁱⁱ	0.91	1.88	2.779 (2)	171

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + 1$; (ii) $x, y - 1, z$; (iii) $-x + 2, y - \frac{1}{2}, -z + 1$.

Data collection: *COLLECT* (Nonius, 1997); cell refinement: *DENZO-SMN* (Otwinowski & Minor, 1997); data reduction: *DENZO-SMN*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL/PC* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL/PC*.

This work was supported by Ningbo Natural Science Fund (grant Nos. 2005B100085).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2591).

References

- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- He, Q., Jennings, M. C., Rohani, S., Zhu, J. & Gomaa, H. (2007). *Acta Cryst. E* **63**, o4199.
- Nonius (1997). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supporting information

Acta Cryst. (2008). E64, o559 [doi:10.1107/S1600536808003516]

(*R*)-1-Phenylethanaminium (*S*)-4-chloromandelate

Quan He, Michael C. Jennings, Sohrab Rohani, Jesse Zhu and Hassan Gomaa

S1. Comment

In our previous work, phenylethylamine (PEA) has been proven to be an efficient resolving agent for resolution of racemic 4-chloromandelic acid. In order to further investigate the chiral recognition mechanism, the single-crystal structure of the corresponding more soluble salt, (*R*)-Phenylethylammonium (*S*)-4-chloromandelate, is reported here for comparison with that of the less soluble salt (*R*)-Phenylethylammonium (*R*)-4-chloromandelate (He *et al.*, 2007).


The title complex consists of an ion pair; an amine cation and a carboxylate anion (see Fig. 1). The absolute stereochemistry of each ion has been confirmed by the structure determination [absolute structure parameter -0.03 (8); (Flack, 1983)]. All three H atoms of the $-\text{NH}_3$ group and the H atom of the O—H group act as hydrogen bond donors in intermolecular O—H···O and N—H···O hydrogen bonds, forming a two-dimensional network perpendicular to the *c* axis. The hydrogen bond motif in the title compound is different to that observed in the room temperature structure of the *R,R* diastereomer (He *et al.*, 2007).

S2. Experimental

To a solution (*S*)-4-chloromandelate (2.0 g, 0.01 mol) in 10 ml 2-propanol, (1.3 mL, 0.01 mol), (*R*)-Phenylethylammonium, was added gradually. A white crystalline solid appeared. The crystals were collected and washed with 2-propanol twice to give the title compound (1.85 g), yield 56%. Single crystals were grown from a concentrated methanol solution of the title compound by slow evaporation at room temperature.

S3. Refinement

All H atoms were positioned geometrically and constrained as riding atoms with; C—H = 1.00 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for methyne H atoms, C—H = 0.98 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H atoms, C—H = 0.95 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic H atoms, O—H = 0.84 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for hydroxyl H atoms and N—H = 0.91 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for amine H atoms.

**Figure 1**

The molecular structure of the title compound with 30% probability displacement ellipsoids and the atom labelling scheme.

(R)-1-Phenylethanaminium (S)-4-chloromandelate

Crystal data



$M_r = 307.76$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 10.4091 (7) \text{ \AA}$

$b = 5.7635 (4) \text{ \AA}$

$c = 13.2544 (10) \text{ \AA}$

$\beta = 96.831 (4)^\circ$

$V = 789.52 (10) \text{ \AA}^3$

$Z = 2$

$F(000) = 324$

$D_x = 1.295 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 11634 reflections

$\theta = 2.0\text{--}27.5^\circ$

$\mu = 0.25 \text{ mm}^{-1}$

$T = 150 \text{ K}$

Plate, colourless

$0.45 \times 0.15 \times 0.08 \text{ mm}$

Data collection

Nonius KappaCCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ scans, and ω scans with κ offsets

Absorption correction: multi-scan
(SORTAV; Blessing, 1995)

$T_{\min} = 0.787$, $T_{\max} = 0.984$

7943 measured reflections

3431 independent reflections

2881 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.072$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.4^\circ$

$h = -13 \rightarrow 13$

$k = -7 \rightarrow 7$

$l = -16 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.115$

$S = 1.05$

3431 reflections

192 parameters

1 restraint

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0565P)^2 + 0.1172P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.26 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.30 \text{ e \AA}^{-3}$

Absolute structure: Flack (1983), 1436 Fridel pairs

Absolute structure parameter: -0.03 (8)

Special details

Experimental. Absorption correction: multi-scan from symmetry-related measurements (*SORTAV*; Blessing, 1995). *M.p.* 140.8–142.5 K. The specific rotation was $[\alpha]^{25}_{\text{D}} = +50.5^\circ$ ($c=1$, $\text{C}^1\text{H}^3\text{OH}$), determined using a Perkin Elmer Model 341 Digital Polarimeter; $^1\text{H-NMR}$ (d_6 -DMSO/TMS): δ 1.424 (d, 3H, CH_3), 4.300 (m, 1H, CHNH_2), 4.536 (s, 1H, CHOH), 7.268–7.457 (m, 9H, C_6H_5 and $\text{C}_6\text{H}_4\text{Cl}$) measured using an INOVA 400 MHz NMR (Varian).

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted *R*-factor wR and goodness of fit *S* are based on F^2 , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating *R*-factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.62566 (7)	0.61666 (16)	1.01588 (5)	0.0611 (3)
C2	0.6271 (2)	0.5894 (5)	0.88508 (17)	0.0393 (6)
C3	0.5788 (3)	0.7663 (5)	0.82206 (19)	0.0435 (6)
H3A	0.5408	0.8990	0.8490	0.052*
C4	0.5863 (2)	0.7484 (4)	0.71887 (18)	0.0368 (5)
H4A	0.5529	0.8702	0.6751	0.044*
C5	0.64141 (18)	0.5567 (4)	0.67818 (16)	0.0260 (4)
C6	0.6887 (2)	0.3804 (4)	0.74326 (19)	0.0359 (5)
H6A	0.7267	0.2473	0.7165	0.043*
C7	0.6815 (2)	0.3952 (5)	0.84688 (19)	0.0422 (6)
H7A	0.7137	0.2730	0.8909	0.051*
C8	0.64285 (18)	0.5435 (3)	0.56394 (15)	0.0246 (4)
H8A	0.6574	0.7033	0.5382	0.030*
O9	0.52152 (12)	0.4619 (3)	0.51580 (11)	0.0287 (3)
H9A	0.4701	0.5740	0.5053	0.043*
C10	0.74941 (19)	0.3868 (4)	0.53386 (15)	0.0249 (4)
O11	0.72224 (14)	0.1839 (3)	0.50678 (13)	0.0345 (4)
O12	0.86102 (13)	0.4771 (3)	0.53936 (11)	0.0299 (3)
N13	0.87681 (16)	-0.1239 (3)	0.42117 (14)	0.0269 (4)
H13A	0.8657	-0.2587	0.4548	0.040*
H13B	0.8332	-0.0076	0.4487	0.040*
H13C	0.9625	-0.0885	0.4267	0.040*
C14	0.6876 (2)	-0.2471 (5)	0.30409 (19)	0.0395 (6)
H14A	0.6898	-0.4075	0.3288	0.059*
H14B	0.6480	-0.2430	0.2332	0.059*
H14C	0.6366	-0.1517	0.3458	0.059*
C15	0.8256 (2)	-0.1521 (4)	0.31099 (16)	0.0296 (5)
H15A	0.8223	0.0045	0.2783	0.035*
C16	0.91365 (19)	-0.3052 (4)	0.25701 (17)	0.0278 (5)

C17	0.9388 (2)	-0.2491 (4)	0.15998 (17)	0.0348 (5)
H17A	0.9041	-0.1103	0.1292	0.042*
C18	1.0142 (2)	-0.3926 (5)	0.10662 (18)	0.0425 (6)
H18A	1.0293	-0.3536	0.0394	0.051*
C19	1.0672 (2)	-0.5923 (5)	0.15193 (19)	0.0400 (6)
H19A	1.1194	-0.6903	0.1160	0.048*
C20	1.0443 (2)	-0.6496 (4)	0.24955 (19)	0.0376 (6)
H20A	1.0815	-0.7859	0.2810	0.045*
C21	0.9671 (2)	-0.5080 (4)	0.30114 (17)	0.0325 (5)
H21A	0.9502	-0.5495	0.3677	0.039*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0659 (5)	0.0910 (7)	0.0280 (3)	-0.0131 (4)	0.0120 (3)	-0.0052 (4)
C2	0.0348 (12)	0.0566 (15)	0.0273 (11)	-0.0126 (12)	0.0076 (9)	-0.0033 (12)
C3	0.0474 (15)	0.0465 (15)	0.0389 (14)	0.0067 (12)	0.0141 (11)	-0.0070 (13)
C4	0.0438 (13)	0.0333 (12)	0.0342 (12)	0.0085 (11)	0.0089 (10)	-0.0005 (10)
C5	0.0202 (9)	0.0275 (11)	0.0311 (10)	-0.0025 (8)	0.0058 (8)	-0.0028 (9)
C6	0.0376 (12)	0.0329 (11)	0.0383 (13)	0.0040 (10)	0.0094 (10)	0.0050 (11)
C7	0.0415 (13)	0.0485 (15)	0.0366 (13)	0.0003 (12)	0.0048 (10)	0.0083 (12)
C8	0.0202 (9)	0.0239 (10)	0.0299 (11)	-0.0023 (8)	0.0035 (8)	-0.0004 (8)
O9	0.0199 (7)	0.0278 (8)	0.0376 (9)	0.0015 (6)	0.0000 (6)	-0.0031 (7)
C10	0.0223 (9)	0.0273 (11)	0.0256 (10)	0.0011 (8)	0.0052 (8)	0.0004 (9)
O11	0.0290 (8)	0.0286 (8)	0.0474 (10)	-0.0002 (6)	0.0104 (7)	-0.0071 (7)
O12	0.0208 (7)	0.0323 (8)	0.0373 (8)	-0.0001 (6)	0.0067 (6)	0.0026 (7)
N13	0.0233 (8)	0.0258 (9)	0.0325 (9)	0.0003 (7)	0.0066 (7)	-0.0027 (8)
C14	0.0272 (11)	0.0549 (15)	0.0361 (13)	0.0034 (11)	0.0029 (9)	-0.0099 (12)
C15	0.0290 (11)	0.0321 (12)	0.0284 (11)	0.0066 (9)	0.0063 (9)	0.0001 (9)
C16	0.0226 (10)	0.0298 (11)	0.0314 (11)	-0.0001 (8)	0.0051 (8)	-0.0043 (9)
C17	0.0373 (12)	0.0359 (12)	0.0320 (12)	0.0035 (10)	0.0070 (9)	0.0021 (10)
C18	0.0429 (13)	0.0526 (16)	0.0344 (12)	0.0052 (12)	0.0145 (10)	-0.0043 (12)
C19	0.0304 (11)	0.0468 (15)	0.0442 (14)	0.0054 (10)	0.0105 (10)	-0.0116 (12)
C20	0.0338 (12)	0.0358 (13)	0.0432 (14)	0.0078 (10)	0.0040 (10)	-0.0038 (11)
C21	0.0340 (11)	0.0343 (12)	0.0305 (12)	0.0038 (10)	0.0087 (9)	0.0003 (10)

Geometric parameters (\AA , $^\circ$)

C11—C2	1.743 (2)	N13—H13B	0.9100
C2—C3	1.375 (4)	N13—H13C	0.9100
C2—C7	1.378 (4)	C14—C15	1.530 (3)
C3—C4	1.383 (3)	C14—H14A	0.9800
C3—H3A	0.9500	C14—H14B	0.9800
C4—C5	1.384 (3)	C14—H14C	0.9800
C4—H4A	0.9500	C15—C16	1.512 (3)
C5—C6	1.385 (3)	C15—H15A	1.0000
C5—C8	1.518 (3)	C16—C17	1.381 (3)
C6—C7	1.387 (3)	C16—C21	1.393 (3)

C6—H6A	0.9500	C17—C18	1.390 (3)
C7—H7A	0.9500	C17—H17A	0.9500
C8—O9	1.425 (2)	C18—C19	1.382 (4)
C8—C10	1.520 (3)	C18—H18A	0.9500
C8—H8A	1.0000	C19—C20	1.383 (3)
O9—H9A	0.8400	C19—H19A	0.9500
C10—O11	1.246 (3)	C20—C21	1.382 (3)
C10—O12	1.267 (2)	C20—H20A	0.9500
N13—C15	1.503 (3)	C21—H21A	0.9500
N13—H13A	0.9100		
C3—C2—C7	121.1 (2)	H13A—N13—H13C	109.5
C3—C2—Cl1	119.5 (2)	H13B—N13—H13C	109.5
C7—C2—Cl1	119.4 (2)	C15—C14—H14A	109.5
C2—C3—C4	119.1 (2)	C15—C14—H14B	109.5
C2—C3—H3A	120.5	H14A—C14—H14B	109.5
C4—C3—H3A	120.5	C15—C14—H14C	109.5
C3—C4—C5	121.3 (2)	H14A—C14—H14C	109.5
C3—C4—H4A	119.3	H14B—C14—H14C	109.5
C5—C4—H4A	119.3	N13—C15—C16	110.97 (17)
C4—C5—C6	118.5 (2)	N13—C15—C14	108.59 (18)
C4—C5—C8	118.86 (19)	C16—C15—C14	112.37 (19)
C6—C5—C8	122.64 (19)	N13—C15—H15A	108.3
C5—C6—C7	121.0 (2)	C16—C15—H15A	108.3
C5—C6—H6A	119.5	C14—C15—H15A	108.3
C7—C6—H6A	119.5	C17—C16—C21	118.5 (2)
C2—C7—C6	119.1 (2)	C17—C16—C15	119.8 (2)
C2—C7—H7A	120.4	C21—C16—C15	121.69 (19)
C6—C7—H7A	120.4	C16—C17—C18	121.0 (2)
O9—C8—C5	110.47 (15)	C16—C17—H17A	119.5
O9—C8—C10	108.78 (16)	C18—C17—H17A	119.5
C5—C8—C10	112.68 (16)	C19—C18—C17	119.6 (2)
O9—C8—H8A	108.3	C19—C18—H18A	120.2
C5—C8—H8A	108.3	C17—C18—H18A	120.2
C10—C8—H8A	108.3	C18—C19—C20	120.1 (2)
C8—O9—H9A	109.5	C18—C19—H19A	120.0
O11—C10—O12	125.27 (19)	C20—C19—H19A	120.0
O11—C10—C8	119.05 (17)	C21—C20—C19	119.8 (2)
O12—C10—C8	115.68 (18)	C21—C20—H20A	120.1
C15—N13—H13A	109.5	C19—C20—H20A	120.1
C15—N13—H13B	109.5	C20—C21—C16	121.0 (2)
H13A—N13—H13B	109.5	C20—C21—H21A	119.5
C15—N13—H13C	109.5	C16—C21—H21A	119.5
C7—C2—C3—C4	-0.5 (4)	C5—C8—C10—O11	98.5 (2)
Cl1—C2—C3—C4	176.8 (2)	O9—C8—C10—O12	155.95 (17)
C2—C3—C4—C5	-0.1 (4)	C5—C8—C10—O12	-81.2 (2)
C3—C4—C5—C6	0.4 (3)	N13—C15—C16—C17	-139.6 (2)

C3—C4—C5—C8	178.0 (2)	C14—C15—C16—C17	98.6 (2)
C4—C5—C6—C7	-0.2 (3)	N13—C15—C16—C21	42.8 (3)
C8—C5—C6—C7	-177.7 (2)	C14—C15—C16—C21	-79.0 (3)
C3—C2—C7—C6	0.8 (4)	C21—C16—C17—C18	0.9 (3)
C11—C2—C7—C6	-176.52 (19)	C15—C16—C17—C18	-176.8 (2)
C5—C6—C7—C2	-0.4 (3)	C16—C17—C18—C19	-1.3 (4)
C4—C5—C8—O9	-81.8 (2)	C17—C18—C19—C20	0.5 (4)
C6—C5—C8—O9	95.7 (2)	C18—C19—C20—C21	0.8 (3)
C4—C5—C8—C10	156.32 (19)	C19—C20—C21—C16	-1.2 (3)
C6—C5—C8—C10	-26.2 (3)	C17—C16—C21—C20	0.4 (3)
O9—C8—C10—O11	-24.4 (3)	C15—C16—C21—C20	178.1 (2)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
O9—H9A···O9 ⁱ	0.84	2.26	2.939 (2)	139
O9—H9A···O11 ⁱ	0.84	2.09	2.826 (2)	146
N13—H13A···O12 ⁱⁱ	0.91	1.89	2.798 (2)	172
N13—H13B···O11	0.91	1.83	2.731 (2)	169
N13—H13C···O12 ⁱⁱⁱ	0.91	1.88	2.779 (2)	171

Symmetry codes: (i) $-x+1, y+1/2, -z+1$; (ii) $x, y-1, z$; (iii) $-x+2, y-1/2, -z+1$.